트레드밀 운동이 외상성 뇌손상 흰쥐 해마에서 신경세포생성 및 BDNF 발현에 미치는 영향

이희혁¹⁾ · 윤진환¹⁾ · 정일규¹⁾ · 김종오¹⁾ · 김보균²⁾ · 서진희²⁾ · 고일규²⁾ · 신말순²⁾ · 김성은²⁾ · 김창주²⁾

1) 한남대학교

2) 경희대학교

Abstract

초 록

Lee, H.H., Yoon, J.H., Jeong, I.G., Kim, J.O., Kim, B.K., Seo J.H., Ko, I.G., Shin, M.S., Kim, S.E., Kim, C.J. Effects of treadmill exercise on cell proliferation and BDNF expression in the rat hippocampus after traumatic brain injury. *Exercise Science*, 17(3): 363-372, 2008. In the present study, we investigated the effects of treadmill exercise on memory function and brain-derived neurotrophic factor (BDNF) levels in rat hippocampus following traumatic brain injury. Male SD rats, aged 6 weeks, were randomly assigned to the following three groups: Sham-operation group(n=10), TBI group(n=10), and TBI and exercise group(n=10). Traumatic brain injury(TBI) was induced by a electromagnetic controlled cortical impact. Treadmill running was conducted with duration and frequency of 30 minutes and 7 times per week, respectively, for 2 weeks. Rats were tested on a passive avoidance task in order to evaluate the memory ability. Hippocampal cell proliferation and BDNF expression were assessed by BrdU immunohistochemistry and Western blot, respectively. In our results, TBI resulted in significant memory impairment along with suppressed BDNF expression in the hippocampus. Treadmill exercise significantly improved the memory capability and increased the hippocampal BDNF expression following TBI. In addition, treadmill exercise upregulated TBI-induced cell proliferation in the hippocampus. Based on the present results, it can be suggested that treadmill running exert a beneficial effect on TBI-induced memory impairment partially by enhancing cell proliferation and BDNF expression in the hippocampus.

이희혁, 윤진환, 정일규, 김종오, 김보균, 서진희, 고일규, 신말순, 김성은, 김창주. 트레드밀 운동이 외상성 뇌손상 흰쥐 해마에서 신경세포생성 및 BDNF 발현에 미치는 영향. *운동과학*, 제17권 제3호, 363-372, 2008. 본 연구는 외상성 뇌손상 흰쥐에서 운동의 기억력 향상기전을 해마 BDNF 발현과 함께 해마 신경세포생성을 조사해 검증하는 것이다. 본 연구는 SD계열 수컷 흰쥐 30마리를 대조군(n=10), 외상성 뇌손상군(n=10), 그리고 외상성 뇌손상과 운동군(n=10)으로 분류하였다. 외상성 뇌손상 흰쥐모델은 전자기적 피질 충격(electromagnetic controlled cortical impact)기를 이용해 유도했다. 운동군의 동물들은 트레드밀에서 매일 30분씩 2주간 달리도록 하였다. 기억력 검사는 passive avoidance 수행검사 그리고 신경세포생성과 BDNF 단백질 검사는 BrdU 면역조직 화학검사와 Western blot을 각각 이용하였다. 본 연구 결과 대조군에 비해 외상성 뇌손상군에서 기억력이 유의하게 감소하였으나, 트레드밀 운동은 외상성 뇌손상으로 유발된 기억력 손상을 완화시키는 것으로 나타났다. 또한 외상성 뇌손상 동물에서 해마 BDNF 발현이 감소되었으나, 운동은 BDNF 단백질 발현과 신경세포생성을 유의하게 증가시켰다. 이러한 결과는 운동이 외상성 뇌손상 동물에서 BDNF 단백질을 증가시켜 신경세포생성을 촉진시킴으로써 기억력 손상을 완화시키는데 도움이 될 수 있음을 보여주는 것이다.

Key words : treadmill exercise, traumatic brain injury, hippocampus, cell proliferation, BDNF

주요어 : 트레드밀 운동, 외상성 뇌손상, 해마, 신경세포생성, 신경영양인자

참고문헌

- Anderson, B. J., Rapp, D. N., Baek, D. H., McCloskey, D. P., Coburn-Litvak, P. S., & Robinson, J. K.(2000). Exercise influences spatial learning in the radial arm maze. *Physiology and Behavior*, 70: 425-429.
- Baranova, A. I., Whiting, M. D., & Hamm, R. J.(2006). Delayed, post-injury treatment with aniracetam improves cognitive performance after traumatic brain injury in rats. *J. Neurotrauma*, 23: 1233-1240.
- Blaha, G. R., Raghupathi, R., Saatman, K. E., & McIntosh, T. K.(2000). Brain-derived neurotrophic factor administration after traumatic brain injury in the rat does not protect against behavioral or histological deficits. *Neuroscience*, 99: 483-493.
- Brody, D. L., Mac-Donald, C., Kessens, C. C., Yuede, C., Parsadarian, M., Spinner, M., Kim, E., Schwetye, K. E., Holtzman, D. M., & Bayly, P. V.(2007). Electromagnetic controlled cortical impact device for precise, graded experimental traumatic brain injury. *J. Neurotrauma*, 4: 657-673.
- Carro, E., Trejo, J. L., Busiguina, S., & Torres-Aleman, I.(2001). Circulating insulin-like growth factor I mediates the protective effects of physical exercise against brain insults of different etiology and anatomy. *Journal of Neuroscience*, 21: 5678-5684.
- Cohen, A. S., Pfister, B. J., Schwarzbach, E., Grady, M. S., Goforth, P. B., & Satin, L. S.(2007). Injury-induced alterations in CNS electrophysiology. *Prog. Brain Res.*, 161: 143-169.
- Cotman, C. W., & Berchtold, N. C.(2002). Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends Neurosci.*, 25: 295-301.
- Deweere, B., Pillon, B., Pochon, J. B., & Dubois, B.(2001). Is the HM story only a "remote memory"? Some facts about hippocampus and memory in humans. *Behav. Brain Res.*, 127: 209-224.
- Ding, Y. H., Li, J., Yao, W. X., Rafols, J. A., Clark, J. C., & Ding, Y.(2006). Exercise preconditioning upregulates cerebral integrins and enhances cerebrovascular integrity in ischemic rats. *Acta. Neuropathol.*, 112: 70-84.
- Donovan, M. H., Yamaguchi, M., & Eisch, A. J.(2008). Dynamic expression of TrkB receptor protein on proliferating and maturing cells in the adult mouse dentate gyrus. *Hippocampus*, 18: 435-439.
- Emery, C. F., & Gatz, M.(1990). Psychological and cognitive effects of an exercise program for community-residing older adults. *Gerontologist*, 30: 184-188.
- Farmer, J., Zhao, X., van Praag, H., Wodtke, K., Gage, F. H., & Christie, B. R.(2004). Effects of voluntary exercise on synaptic plasticity and gene expression in the dentate gyrus of adult male Sprague-Dawley rats in vivo. *Neuroscience*, 124: 71-79.
- Flanagan, S. R., Hibbard, M. R., Riordan, B., & Gordon, W. A.(2006). Traumatic brain injury in the elderly: diagnostic and treatment challenges. *Clin. Geriatr. Med.*, 22: 449-468.
- Fordyce, D. E., & Wehner, J. M.(1993). Physical activity enhances spatial learning performance with an associated alteration in hippocampal protein kinase C activity in C57BL/6 and DBA/2 mice. *Brain Res.*, 619: 111-119.
- Gao, J., Prough, D. S., McAdoo, D. J., Grady, J. J., Parsley, M. O., Ma, L., Tarensenko, Y. I., & Wu, P.(2006). Transplantation of primed human fetal neural stem cells improves cognitive function in rats after traumatic brain injury. *Exp. Neurol.*, 201: 281-292.
- Gomez-Pinilla, F., So, V., & Kesslak, J. P.(1998). Spatial learning and physical activity contribute to the induction of fibroblast growth factor: neural substrates for increased cognition associated with exercise. *Neuroscience*, 85: 53-61.
- Graham, D. I., Adams, J. H., Nicoll, J. A., Maxwell, W. L., & Gennarelli, T. A.(1995). The nature, distribution and causes of traumatic brain injury. *Brain Pathol.*, 5: 397-406.
- Griesbach, G. S., Gomez-Pinilla, F., & Hovda, D. A.(2004). The upregulation of plasticity-related proteins following TBI is disrupted with acute voluntary

- exercise. *Brain Res.*, 1016: 154-162.
- Hatta, A., Nishihira, Y., Kim, S. R., Kaneda, T., Kida, T., Kamijo, K., Sasahara, M., & Haga, S.(2005). Effects of habitual moderate exercise on response processing and cognitive processing in older adults. *Jpn. J. Physiol.*, 55: 29-36.
- Hicks, R. R., Keeling, K. L., Yang, M. Y., Smith, S. A., Simons, A. M., & Kotwal, G. J.(2002). Vaccinia virus complement control protein enhances functional recovery after traumatic brain injury. *J. Neurotrauma*, 19: 705-714.
- Johansson, B. B., & Ohlsson, A. L.(1996). Environment, social interaction, and physical activity as determinants of functional outcome after cerebral infarction in the rat. *Exp. Neurol.*, 139: 322-327.
- Kempermann, G., Wiskott, L., & Gage, F. H.(2004). Functional significance of adult neurogenesis. *Curr. Opin. Neurobiol.*, 14: 186-191.
- Kim, Y. P., Kim, H. B., Jang, M. H., Lim, B. V., Kim, Y. J., Kim, H., Kim, S. S., Kim, E. H., & Kim, C. J.(2003). Magnitude-and time-dependence of the effect of treadmill exercise on cell proliferation in the dentate gyrus of rats. *Int. J. Sports Med.*, 24: 114-117.
- Kleindienst, A., McGinn, M. J., Harvey, H. B., Colello, R. J., Hamm, R. J., & Bullock, M. R.(2005). Enhanced hippocampal neurogenesis by intraventricular S100B infusion is associated with improved cognitive recovery after traumatic brain injury. *J. Neurotrauma*, 22: 645-655.
- Kotani, S., Yamauchi, T., Teramoto, T., & Ogura, H.(2006). Pharmacological evidence of cholinergic involvement in adult hippocampal neurogenesis in rats. *Neuroscience*, 142, 505-514.
- Larner, S. F., McKinsey, D. M., Hayes, R. L., & Wang, K. K.(2005). Caspase 7: increased expression and activation after traumatic brain injury in rats. *J. Neurochem.*, 94: 97-108.
- Larson, E. B., Wang, L., Bowen, J. D., McCormick, W. C., Teri, L., Crane, P., & Kukull, W.(2006). Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. *Ann. Intern. Med.*, 144: 73-81.
- Lie, D. C., Song, H., Colamarino, S. A., Ming, G. L., & Gage, F. H.(2004). Neurogenesis in the adult brain: new strategies for central nervous system diseases. *Annu. Rev. Pharmacol. Toxicol.*, 44: 399-421.
- Lu, D., Mahmood, A., Qu, C., Goussev, A., Schallert, T., & Chopp, M.(2005). Erythropoietin enhances neurogenesis and restores spatial memory in rats after traumatic brain injury. *J. Neurotrauma*, 22: 1011-1017.
- Lu, D., Qu, C., Goussev, A., Jiang, H., Lu, C., Schallert, T., Mahmood, A., Chen, J., Li, Y., & Chopp, M.(2007). Statins increase neurogenesis in the dentate gyrus, reduce delayed neuronal death in the hippocampal CA3 region, and improve spatial learning in rat after traumatic brain injury. *J. Neurotrauma*, 24: 1132-1146.
- Mahmood, A., Lu, D., Qu, C., Goussev, A., Zhang, Z. G., Lu, C., & Chopp, M.(2007). Treatment of traumatic brain injury in rats with erythropoietin and carbamylated erythropoietin. *J. Neurosurg*, 107: 392-397.
- Mizuno, M., Yamada, K., Olariu, A., Nawa, H., & Nabeshima, T.(2000). Involvement of brain-derived neurotrophic factor in spatial memory formation and maintenance in a radial arm maze test in rats. *J. Neurosci.*, 20: 7116-7121.
- Narayan, R. K., Michel, M. E., Ansell, B., Baethmann, A., Biegon, A., & Bracken, M. B.(2002). Clinical trials in head injury. *J. Neurotrauma*, 19: 503-557.
- Sairanen, M., Lucas, G., Ernfors, P., Castrén, M., & Castrén, E.(2005). Brain-derived neurotrophic factor and antidepressant drugs have different but coordinated effects on neuronal turnover, proliferation, and survival in the adult dentate gyrus. *J. Neurosci.*, 25: 1089-1094.
- Schmidt, R. H., Scholten, K. J., & Maughan, P. H.(1999). Time course for recovery of water maze performance and central cholinergic innervation after fluid percussion injury. *J. Neurotrauma*,

16: 1139-1147.

- Shors, T. J., Miesegaes, G., Beylin, A., Zhao, M., Rydel, T., & Gould, E.(2001). Neurogenesis in the adult is involved in the formation of trace memories. *Nature*, 410: 372-376.
- Smith, D. H., Okiyama, K., Thomas, M. J., Claussen, B., & McIntosh, T. K.(1991). Evaluation of memory dysfunction following experimental brain injury using the Morris water maze. *J. Neurotrauma*, 8: 259-269.
- Sugaya, K., Alvarez, A., Marutle, A., Kwak, Y. D., & Choumkin, E.(2006). Stem cell strategies for Alzheimer's disease therapy. *Panminerva. Med.*, 48: 87-96.
- Sun, D., McGinn, M. J., Zhou, Z., Harvey, H. B., Bullock, M. R., & Colello, R. J.(2007). Anatomical integration of newly generated dentate granule neurons following traumatic brain injury in adult rats and its association to cognitive recovery. *Exp. Neurol.*, 204: 264-72.
- Swain, R. A., Harris, A. B., Wiener, E. C., Dutka, M. V., Morris, H. D., Theien, B. E., Konda, S., Engberg, K., Lauterbur, P. C., & Greenough, W. T.(2003). Prolonged exercise induces angiogenesis and increases cerebral blood volume in primary motor cortex of the rat. *Neuroscience*, 117: 1037-1046.
- Trejo, J. L., Carro, E., & Torres-Aleman, I.(2001). Circulating insulin-like growth factor I mediates exercise-induced increases in the number of new neurons in the adult hippocampus. *J. Neurosci.*, 21: 1628-1634.
- van Praag, H., Kempermann, G., & Gage, F. H.(1999). Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus. *Nat. Neurosci.*, 2: 266-270.
- Vaynman, S., Ying, Z., & Gomez-Pinilla, F.(2004). Hippocampal BDNF mediates the efficacy of exercise on synaptic plasticity and cognition. *Eur. J. Neurosci.*, 20: 2580-2590.
- Wagner, A. K., Chen, X., Kline, A. E., Li, Y., Zafonte, R. D., & Dixon, C. E.(2005). Gender and environmental enrichment impact dopamine transporter expression after experimental traumatic brain injury. *Exp. Neurol.*, 195: 475-483.
- Witgen, B. M., Lifshitz, J., Smith, M. L., Schwarzbach, E., Liang, S. L., Grady, M. S., & Cohen, A. S.(2005). Regional hippocampal alteration associated with cognitive deficit following experimental brain injury: a systems, network and cellular evaluation. *Neuroscience*, 133: 1-15.
- Wu, H., Lu, D., Jiang, H., Xiong, Y., Qu, C., Li, B., Mahmood, A., Zhou, D., & Chopp, M.(2008). Simvastatin-mediated upregulation of VEGF and BDNF, activation of the PI3K/Akt pathway, and increase of neurogenesis are associated with therapeutic improvement after traumatic brain injury. *J. Neurotrauma*, 25: 130-139.
- Xiong, Y., Mahmood, A., Lu, D., Qu, C., Goussev, A., Schallert, T., & Chopp, M.(2007). Role of gender in outcome after traumatic brain injury and therapeutic effect of erythropoietin in mice. *Brain Res.*, 1185: 301-312.
- Yasuhara, T., Matsukawa, N., Hara, K., Yu, G., Xu, L., Maki, M., Kim, S. U., & Borlongan, C. V.(2006). Transplantation of human neural stem cells exerts neuroprotection in a rat model of Parkinson's disease. *J. Neurosci.*, 26: 12497-12511.

논문투고일 : 2008. 06. 09

심사일 : 2008. 07. 14

심사완료일 : 2008. 08. 12